


289

FSTC-HT-23-512-68

# U.S. ARMY FOREIGN SCIENCE AND TECHNOLOGY CENTER



AD 685449


PRODUCTION OF A LETHAL TOXIN AND CL. HISTOLYTICUM

COLLAGENASE ON CASEIN NUTRITIVE MEDIA

COUNTRY: USSR

## TECHNICAL TRANSLATION

Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.



TECHNICAL TRANSLATION

FSTC-HT-23- 512-68

PRODUCTION OF A LETHAL TOXIN AND CL. HISTOLYTICUM  
COLLAGENASE ON CASEIN NUTRITIVE MEDIA

by

Ye. V. Vlasova and F. F. Tsurikov

SOURCE: ZHURNAL MIKROBIOLOGII, EPIDEMIOLOGII I IMMUNOBIOLOGII  
No. 1, pp. 112-114, 1966  
USSR

Translated for FSTC by Techtran Corporation

This translation is a rendition of the original foreign text without any analytical or editorial comment. Statements or theories advocated or implied are those of the source and do not necessarily reflect the position or opinion of the US Army Foreign Science and Technology Center. This translation is published with a minimum of copy editing and graphics preparation in order to expedite the dissemination of information. Requests for additional copies of this document should be addressed to the Defense Documentation Center, Cameron Station, Alexandria, Virginia, ATTN: OSR-2

UNCLASSIFIED  
Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)

Foreign Science and Technology Center  
US Army Materiel Command  
Department of the Army

2a. REPORT SECURITY CLASSIFICATION

UNCLASSIFIED

2b. GROUP

3. REPORT TITLE

Production of a Lethal Toxin and Cl. Histolyticum Collagenase on Casein  
Nutritive Media

4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

Translation

5. AUTHOR(S) (First name, middle initial, last name)

Ye. V. Vlasova and F. F. Tsurikov

6. REPORT DATE

FEB 18 1969

7a. TOTAL NO. OF PAGES

6

7b. NO. OF REFS

N/A

8a. CONTRACT OR GRANT NO.

a. PROJECT NO. 9803601020300

c. 9223628 2301

d.

9a. ORIGINATOR'S REPORT NUMBER(S)

ESTC-HT-23-512-68

OTHER REPORT NO(S) (Any other numbers that may be assigned  
this report)

ACSI Control Number (None)

10. DISTRIBUTION STATEMENT

Distribution of this document is unlimited.

11. SUPPLEMENTARY NOTES

12. SPONSORING MILITARY ACTIVITY

US Army Foreign Science and Technology  
Center

13. ABSTRACT

Three strains of Cl. histolyticum were grown on various nutritive media, and a-toxin and collagenase were extracted, purified, and concentrated from culture filtrates. The a-toxin was present at levels of 500-1,000 Dlm/ml, along with 0.5-2 units of collagenase (as indicated by viscometry). Data are presented on the dynamics of a a-toxin and collagenase formation on the suggested nutritive media.

Best Available Copy

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS  
OBSOLETE FOR ARMY USE.

UNCLASSIFIED  
Security Classification

UNCLASSIFIED  
Security Classification

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	W	ROLE	WT
	Cl. histolyticum a-toxin Collagenase Casein nutritive medium						

UNCLASSIFIED  
Security Classification

PRODUCTION OF A LETHAL TOXIN AND *CL. HISTOLYTICUM*  
COLLAGENASE ON CASEIN NUTRITIVE MEDIA

*Cl. histolyticum* is one of the inducers of gaseous gangrene and, although the disease is encountered in a small percent of cases, one must nevertheless consider its influence during construction of immunogenic associated preparations for the prophylaxis of gaseous gangrene.

Interest in this microbe has not been exhausted. It is one of the few microorganisms which produces the enzyme collagenase. Moreover, it is one of the most active producers of collagenase, which is the only enzyme which cleaves native collagen.

In recent years there have been a great number of reports, connected with extraction of collagenase, its purification and concentration, and also its application to the study of the structure of collagen. There have also been reports on the application of collagenase with a medical purpose--for the resolution of keloid projections, treatment of burns, etc. The literature on these questions is already so broad, that we have allowed ourselves to cite only the available reviews of Kunina and Shpikiter (1960) and Mandl (1961).

In agreement with the data of Oakley and Warrak (1950), a number of antigenic components are observed in the filtrate of *Cl. histolyticum* cultures, which they designated  $\alpha$ ,  $\beta$ , and  $\gamma$ ; of these,  $\alpha$  is a lethal toxin, which appears to also have the necrotic effect of  $\beta$ -collagenase and  $\gamma$ -cystein-activating protease. Antibodies towards all three components were detected in the serum of immunized animals. The roles of each of these components in infection and immunity has not at this time been determined, since they have not been obtained in pure form.

Almeier (1952) has shown that immunization with anacollagenase (of which the method of extraction has not been specified) insures better protection from infection than immunization with anatoxin.

From all that has been stated above, it is clear that it is of theoretical and practical interest to obtain filtrates of *Cl. histolyticum*, containing maximal quantities of the  $\alpha$ ,  $\beta$ , and  $\gamma$  components. The aim of the present work was also the development of a recipe for nutritive medium and conditions of cultivation of *Cl. histolyticum*, in order to obtain maximally active filtrates, containing  $\alpha$ -toxin and collagenase, which would later serve as the starting material for extracting purified and concentrated toxins and anatoxins and collagenase.

Four strains of *Cl. histolyticum*--No. 5, 247, H<sub>4</sub> and 230-2--were taken for the work. The strains were kept in nutritive medium from Poup broth with small pieces of fresh meat, 0.1% agar, and 0.4% gelatine under liquid petrolatum.

The toxin was titrated by the generally accepted method in mice by intravenous injection.

The activity of the collagenase was determined by a viscometric method.<sup>1</sup> The activity was marked under conditions of unit-cotangent angles of a straight slope, obtained by means of division of the difference in quantities of the logarithms of the specific viscosity by the difference in the corresponding time intervals in minutes (Kazakova et al., 1958).


Acid, pancreatic hydrolysates of casein and a hydrolysate, obtained with the help of *Aspergillus terricola* protease (Vinogradova, et al., 1958) served as a basis for obtaining various nutritive media. To this hydrolysate were added gross factors--various concentrations of corn extract, yeast water, vitamin B complex (Vinogradova, et al., 1963).

Preliminary data indicated that the best results in relation to production both of  $\alpha$ -toxins, and of collagenase, were obtained in nutritive media based on the acid hydrolysate of casein. Precisely these media were used for the preparation of large quantities of filtrates for the purpose indicated above. Inoculations were carried out in a bottle with cotton in a 2.5 l of nutritive medium and remained in a thermostat at 37°.

---

<sup>1</sup>The determination of collagenase activity was carried out by colleagues at the Institute of Biological and Medical Chemistry by G. A. Levnikov and N. I. Solov'ev, and also a colleague of the Department of Wound Infections of the Gamalei Institute of Epidemiology and Microbiology, G. F. Shemanov.

In the first experiments, samples were taken from the bottles 24, 48, 72, and 96 hours after inoculation, with the purpose of determining the time of maximal accumulation of  $\alpha$ -toxins and collagenase.



Accumulation of  $\alpha$ -Toxin (1, 2) and Collagenase (3, 4) in Two Experiments.

The maximal accumulation of  $\alpha$ -toxins was observed 24 hours, and of collagenase--48 hours after inoculation, while the accumulation of these components occurred independently of each other (cf. Figure). In order to obtain anatoxins, 0.4% formalin was added to the filtrate of the 24-hour culture, and placed in a thermostat at 37° to render it harmless.

In order to obtain purified collagenase, the filtrate of the 48-hour culture was taken. In 1962 we had taken the additional strains of *Cl. histolyticum*, H<sub>4</sub> and 230-2. According to the data of MacLennan et al. (1953) and Mandl and Zaffuto (1960), these strains are the best producers of collagenase. Moreover, addition of 5% peptone to the casein nutritive medium seems to be foolproof for the production of collagenase. These data are confirmed in our preliminary experiments.

In our application of these media without addition of peptone, the growth of the microbes was weak, as was the production of  $\alpha$ -toxin and collagenase. Upon addition of 3 to 5% peptone to the medium, good results were observed in relation both to production of  $\alpha$ -toxin, and collagenase, while the strain *Cl. histolyticum* H<sub>4</sub> gave the worst results in comparison with the strain 230-2, and therefore the latter was taken for further work.

As can be seen from the preliminary data in the table, in our use of casein nutritive media toxin was obtained with a strength of 500 to 1,000 DZm/mZ, i.e. according to activity, they did not yield the toxins, which were obtained by domestic and foreign authors in meat nutritive media (Zyelyevinskaya, 1943; Stewart, 1936; Walbum and Raymann, 1938; Bowen, 1952; Raynand et al., 1954; Wildfuhr, 1950, etc.).

Unfortunately, we cannot compare our data on the activity of collagenase with the data of other authors, because the determination of collagenase activity was carried by different methods. However, the possibility of obtaining sufficient quantities of partially purified collagenase from these filtrates (Vlasova and Solov'eva, 1962) already speaks about collagenase activity obtained from filtrates.

Extraction of Toxin and Collagenase During Cultivation of  
*Cl. histolyticum* in Casein Nutritive Media

Nutritive Medium	Strain	Number of Inno- culations With Various Strains of Toxin (in DZm)			Number of Innoculations with Various Collagenase Activities		
		200-500	500-1000	1000-2000	0.25-0.5	0.5-1	1-2
Acid Hydrolysate of Casein and 3% Corn Extract	5	3	9	1	Not Determined		
Acid Hydrolysate of Casein and 5% Corn Extract		10	14	8	1	3	14
Acid Hydrolysate of Casein and 0.3-0.5% Vitamin B (Complex)	247	—	3	—	—	—	3
Acid Hydrolysate of Casein, 5% Yeast Water and 3-5% Peptone		—	3	7	—	3	4
Acid Hydrolysate of Casein, 5% Yeast Water and 3-5% Peptone	230-2		6	4		3	4
Acid Hydrolysate of Casein, 0.5% Vitamin B (Complex) and 3% Peptone			5				5



## Conclusions

1. The nutritive media and conditions of culturing of *Cl. histolyticum* which we used, permit one to obtain filtrates of the culture, which are active in relation both to  $\alpha$ -toxin, and collagenase.
2. The filtrates which we obtain served as the starting raw material for obtaining purified and concentrated anatoxin of *Cl. histolyticum* for active immunization, and also as the starting raw material for obtaining purified collagenase.

## REFERENCES

1. Vinogradova, I. N., V. A. Pyetryenko, F. F. Tsurikov, et al., *Material' po Obmenu Op'tom Ministerstva Zdravookhraneniya* [Metabolic Materials with the Experiment of the Ministry of Public Health of the USSR], No. 1/53, p. 32, Moscow, 1957.
2. Vlasova, Ye. V. and N. I. Solov'eva, *Vopr. myed. Khimii*, No. 4, p. 424, 1962.
3. Zyelyevinskaya, S. A., *Rukovodstvo Po S'vorotoshnomu i Vaktsinnomu Delu* [Handbook in Serum and Vaccine Matters], Moscow, p. 284, 1943.
4. Kazakova, O. V., V. N. Oryekhovich and V. O. Shpikiter, *Dokl. AN SSSR*, Vol. 120, No. 2, p. 359, 1958.
5. Kunina, O. V. and V. O. Shpikiter, *Uspeyehi sour. biol.*, Vol. 50, No. 3, p. 294, 1960.
6. Altmeier, W. A. and R. Coith, et al., *Arch. Surg.*, Vol. 65, p. 633, 1952.
7. Bowen, H. E., *Yale J. Biol. Med.*, Vol. 25, p. 124, 1952.
8. Howes, E. L., J. Mandl, and S. Zaffuto, *J. Bact.*, Vol. 79, p. 191, 1960.
9. MacLennan, J. D., J. Mandl and E. L. Howes, *J. Clin. Invest.*, Vol. 32, p. 1317, 1953.

10. Mandl, J., *Advanc. Enzymology*, Vol. 23, p. 164, 1961.
11. Oakley, C. and G. Warrak, *J. gen. Microbiol.*, Vol. 4, p. 365, 1950.
12. Raynand, M. and A. Turpen, et al., *Ann. Inst. Pasteur*, Vol. 88, p. 24, 1955.
13. Stewart, S. E., *Publ. Hlth Rep.*, Vol. 51, p. 1272, 1936..
14. Walbum, L. E. and G. S. Reymann, *J. Path. Bact.*, Vol. 46, p. 315, 1938.
15. Wildfuhr, G., *Z. Immun.--Forsch.*, No. 106, p. 369, 1949.